

Spectra and evolution of two X-ray sources in M81

V. La Parola, G. Peres

DSFA - Sez. di Astronomia, Piazza del Parlamento 1, 90134 Palermo, Italy

G.Fabbiano, D. W. Kim

Harvard Smithsonian Center for Astrophysics, 60 Garden st, Cambridge, MA 02138, USA

Abstract. We analyzed the spectral and temporal features of X-ray sources in M81 using data from all the relevant ROSAT-PSPC/HRI observations. We discuss the main features of the point-like nucleus and of the second brightest source in the field (X-9), whose identification is still unclear.

M81 nuclear source

M81 has a LINER nucleus, consistent with an X-ray point-like source (Dyson, priv. comm.). We show its lightcurve over the ~ 7 years monitored by ROSAT (Fig. 1): each point is one observation (La Parola et al., in prep.). We also looked for spectral variability through hardness ratio (HR) analysis and found that the source has three spectral states: we analysed the three spectra derived adding observations with similar HR. The best spectral fit for the hard phase ($HR > 0.21$) is a two-T Raymond-Smith thermal model ($kT_1 = 0.23 \pm 0.01$ KeV, $kT_2 = 2.4 \pm 0.4$ keV); we can model the two softer phases with a power law + Raymond-Smith with compatible values ($\Gamma \sim 2.1$, $kT \sim 0.4$ keV). Luminosity is $\sim 2 \times 10^{40}$ ergs/sec.

X-9

X-9 is a very bright X-ray source 12.5 arcmin away from M81 nucleus. Its optical counterpart has not been clearly identified yet, and the nature of its emission is still unclear. X-9 is strongly variable, as evident from the light curve in Fig 2. Its rapid variability suggests that the bulk of the emission comes from a point-like source, even if from HRI images we found some hint at a diffuse emission. The spectrum appears to be very complex and variable. We divided all data into two groups, or spectral phases, according to the hardness ratio values of each observation, analysing them separately. We found that the best fit for both groups is a multi-temperature black body disk (successfully used to model galactic black hole candidate- Makishima et al., 2000). The temperature we find for the two data sets are only marginally comatible ($kT_1 = 0.58 \pm 0.07$ keV for the soft phase, $kT_2 = 0.68 \pm 0.04$ keV for the hard phase), so confirming the spectral variability. We also see a soft excess in the spectrum, a possible signature of an extended component.




Figure 1. Light curve of M81 nucleus. Each point is the average flux in one observation: stars - PSPC, diamonds - HRI, circle - SAX. All the instruments agree well and strong variability is present on short (days) and on long (years) time scales.

This work was supported in part by NASA grant NAG5-2946 and NASA contract NAS8-39073(CXC) and in part by MURST. This research has made use of the HEASARC online database and of the ESO online DSS.




Figure 2. Light curve of X-9. Each point is the average flux in one observation. Same notation as in Fig. 1. Also this source shows variability on several time scales.